62	L/957	7	*				1.44	K W
	LIBRAI EGELV FEB11 1957	ED						
PRICE LIST NO. 18	U. S. Department of Agr						12.5	တဲ့
NORDAN'S GRASS FARM "Highest Quality of Superior Turf" Phone 2-5842 or 2-5922 Abbeville, Alabama	The second secon	ςį.	sq. yds.—3.	sq. yds.—4.) sq. yds.—5.	200 sq. yds.—6.	400 sq. yds.—7.	Per yard, 401 sq. yds. or more—8
This price list cancels all previous quotations. Prices are subject to change without notice.	prepaid in o sprig 50	2 to 10 sq. yds	11 to 25	26 to 50	51 to 100	101 to	201 to	401 sq. y
Our superior turf is sold in 12" rug mat form by the square yard. Our inferior turf such as edgings and thin turf is sold by the bushel only.	Samples prepaid Enough to sprig	Per yard,						
SOLID SOD LESS SOIL, F.O.B. FARM								
GRADE "A", ***(note below) Bermuda, Tifgreen (Tifton 320) Bermuda, Tiffine (Tifton 127) Bermuda, Tiflawn (Tifton 57) Bermuda, Sunturf Zoysia, Emerald Zoysia, Meyer (Z-52)	\$5.50 5.50 5.50 5.50 5.50 5.50 \$5.25	6.00 5.50 5.50 6.00 7.00 6.00 4.00	5.75 5.25 5.25 5.75 6.75 5.75 3.75	5.50 5.00 5.00 5.50 6.50 5.50 3.50	5.25 4.75 4.75 5.25 6.25 5.25 3.25	5.00 4.50 4.50 5.00 6.00 5.00 3.00	4.75 4.25 4.25 4.75 5.75 4.75 2.75	4.50 4.00 4.00 4.50 5.50 4.50 2.50
GRADE "B", **(note below)								
Bermuda, Everglades No. 1 Bermuda, Gene Tift Bermuda, Tifgreen (Tifton 328) Bermuda, Tiffine (Tifton 127) Bermuda, Tiflawn (Tifton 57) Bermuda, Sunturf Bermuda, U-3	5.25 5.25 5.25 5.25 5.25 5.25 5.25 5.25	4.00 5.00 4.50 4.50 4.50 5.00 4.00	3.75 4.75 4.25 4.25 4.25 4.75 3.75	3.50 4.50 4.00 4.00 4.00 4.50 3.50	3.25 4.25 3.75 3.75 3.75 4.25 3.25	3.00 4.00 3.50 3.50 3.50 4.00 3.00	2.75 3.75 3.25 3.25 3.25 3.75 2.75	2.50 3.50 3.00 3.00 3.50 2.50
Zoysia, Emerald Zoysia, Matrella Zoysia, Meyer (Z-52)	5.25 5.25 5.25 5.25	6.00 3.75 5.00	5.75 3.50 4.75	5.50 3.25 4.50	5.25 3.00 4.25	5.00 5.00 2.75 4.00	4.75 2.50 3.75	4.50 2.25 3.50
GRADE "C" *(note below) (Solid sod with or with	th without soil)							
Bermuda, Everglades No. 1 Bermuda, Gene Tift Bermuda, Tifgreen (Tifton 328)	\$5.00 5.00 5.00	3.00 3.00 4.00	2.75 2.75 3.75	2.50 2.50 3.50	2.25 2.25 3.25	2.00 2.00 3.00	1.75 1.75 2.75	1.50 1.50 2.50

5.00

5.00

5.00

5.00

5.00

5.00

5.00

5.00

5.00

3.50

3.50

4.00

3.00

3.00

3.00

5.00

3.25

4.00

3.25

3.25

3.75

2.75

2.75

2.75

4.75

3.00

3.75

3.00

3.00

3.50

2.50

2.50

2.50

4.50

2.75

3.50

2.75

2.75

3.25

2.25

2.25

2.25

4.25

2.50

3.25

2.50

2.50

3.00

2.00

2.00

2.00

4.00

2.25

3.00

2.25

2.25

2.75

1.75

1.75

1.75

3.75

2.00

2.75

2.00

2.00

2.50

1.50

1.50

3.50

1.75

2.50

1.50

Bermuda, Tiffine (Tifton 127)

Bermuda, Tiflawn (Tifton 57)

Bermuda, Sunturf

Centipede, Chinese

Zoysia, Meyer (2-52)

Zoysia, Emerald

Zoysia, Matrella

Bermuda, U-3

St. Augustine

CONTACT US FOR THE NEW SUPERIOR TURF GRASSES AS THEY ARE RELEASED.

^{***}Denotes highest quality possible. Certified by Alabama and U. S. Crop Improvement Assns. for purity of strain. Certified by U. S. and Alabama Departments of Agriculture. Grown under irrigation upon fumigated soils and dieldrin treated.

^{**}Certified by U. S. and Alabama Departments of Agriculture. Grown under irrigation upon fumigated soils & dieldrin treated.

^{*}Certified by U. S. and Alabama Departments of Agriculture. Grown under irrigation.

[&]quot;THE NATION'S LARGEST COMMERCIAL ASSORTMENT OF PERMANENT LAWN GRASSES"

Inquire regarding the following: BERMUDAS—South African, Bayshore, Everglades No. 3, Ormond, Florida No. 3, Florida No. 4, Florida Special, Transvaal., Tex-Turf, Ideal, Uganda and Abbie; ZOYSIAS—Common japonica, Yellow Stem japonica, Mascarene, Flagstone, King matrella, Asiatic matrella, Nordan matrella, Mowless matrella, Compacto matrella, Manilagrass, No. 13521 matrella; CENTIPEDES—P. I. & Red Stem; ST. AUGUSTINE—Bitter-Blue Stem; DICHONDRA (ground cover); Mondo japonicus, Monkey Grass, & Lireope. (Seed of Common Zoysia japonica and Centipede only.)

	and the Louisian Control of the Cont	-	0	A.	-
PLUGS, ROUND 2"x2" (SOD & SOIL) F.O.B. FARM	1.	2.	3.	4.	5.
PRICED PER 100; Quantities up to	1000	2000	3000	4000	5000 & more
GRADE "C"					
Bermuda, Tifgreen (Tifton 328)*-**	\$ 5.00	4.50	4.00	3.50	3.25
Bermuda, Tiffine (Tifton 127)*-**	5.00	4.50	4.00	3.50	3.25
Bermuda, Tiflawn (Tifton 57)*-**	5.00	4.50	4.00	3.50	3.25
Zoysia, matrella**	4.50	4.00	3.50	3.00	2.75
Zoysia, Emerald*-**	15.00	14.00	13.00	12.00	11.00
Zoysia, Meyer (Z-52)*-**	5.00	4.50	4.00	3.50	3.00
*Denotes available in Grade "A" with fumigated soil at 20c per 100 mg	ore than a	bove pr	ices.		
11 1 C 1 (D) 11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1					

^{**}Denotes available in Grade "B" with fumigated soil at 10c per 100 more than above prices.

TURF SETTING ESTIMATES

- 1 square yard equals approximately 1 to 2 bushels of loose packed turf, depending on the height of cut at time of digging.
- 1 square yard of the Zoysias will average sprigging 1000 square feet of area.
- 1 square yard of the Bermudas will average sprigging 1000 square feet of area.
- 1 square yard of the Centipedes will average sprigging 500 square feet of area.
- 1 square yard of the St. Augustines will average sprigging 500 square feet of area.
- "Solid Sod Less Soil" will weigh from 7 to 15 pounds per square yard. Weight with soil, 60 to 80 lbs.
- "Solid Sod Less Soil" will displace one cubic foot of packing space per square yard.

YOUR CUSTOMERS SEEKING PERSONAL ADVICE WILL RECEIVE SAME PROMPTLY IF WITH THEIR QUESTIONS THEY WILL ENCLOSE ONE DOLLAR TO COVER TIME, POSTAGE AND SUPPLIES.

TERMS AND CONDITIONS OF SALE

Payment by cash, firm's check, cashier's check or money order payable to NORDAN'S GRASS FARMS must accompany all orders. "Pick up" orders must be placed and paid for at least three days prior to scheduled digging. Orders will be filled on specified dates with weather permitting. WE DO NOT SHIP C.O.D. OR OPEN ACCOUNT.

Our sods are machine cut to convenient sizes for ease of handling. We exercise every care to see that our products are handled in a manner that will keep them fresh and alive. We cannot guarantee that they will either live or grow after they leave our hands. Any plants which prove something other than what they were sold for will be replaced, but in no case will a greater price be allowed for any product replaced than was originally paid.

All quotations are for immediate acceptance. No liability to us should stock be sold upon receipt of order or if injured from heat, cold or neglect in transit, or should strikes or other causes prevent delivery.

F.O.B. SHIPMENTS VIA RAILWAY EXPRESS UNLESS OTHERWISE SPECIFIED.

ORDER FORM:				
NAME			DATE	
ST. & NO.	CITY	Z	ONE	STATE
ENCLOSED FIND \$		SHIP THE FOI		() () Parcel Post Air Mail
SPRIGS OF	,		PLUGS OF	
QUANTITY	KIND OF GRASS	QUANTITY		KIND OF GRASS
QUANTITY SQUARE YARDS	OF KIND OF GRASS	() WITH SOIL	() WITHOUT SOII	() L GRADE
WOULD LIKE DELIVERY ABOUT				

NORDAN'S GRASS FARMS

"For Highest Quality of Superior Turf"

Abbeville, Alabama

EMERALD ZOYSIA

"The Modern Miracle Lawn Grass"

Emerald Zoysia is the first released hybrid between a course textured Japanese Zoysia and a very fine textured Mascarene Zoysia. This offspring has proven superior to all other Zoysia grasses tested for turf quality. Since Emerald is a hybrid from a wide cross it is not grown from seed, but is reproduced vegetatively from sod, sprigs or plugs. The only assurance of getting genuine Emerald Zoysia is by purchasing certified stock. Most outstanding advantages of Emerald Zoysia are:

- 1. The turf quality is unexcelled by any of our other lawn grasses. Emerald is a fine-leafed, dense growing, dark green grass that is thrillingly spongy to walk upon.
- 2. A mature Emerald lawn is so thick and dense that weeds haven't a chance to sprout.
- Emerald grows so slowly that it is no problem to keep out of flower beds, walkways and gardens.
- 4. It is possible to produce a covered lawn area in three months when planted on six inch centers.
- 5. Emerald is shade tolerant even in the Deep South where it may be found growing right up to the base of large trees.
- 6. Emerald Zoysia is more frost-tolerant than any of our other lawn grasses, therefore provides a green lawn for a much longer season. It stays green the year around in some southern parts of the nation.
- 7. Emerald turf is highly resistant to wear.
- 8. The upright growth of the leaves is very slow, therefore it requires less frequent mowings.
- 9. No serious damage from diseases or insects has been observed on Emerald Zoysia.
- 10. Emerald Zoysia produces such a dense turf that it maintains its erect body throughout the winter and lends itself to overseeding with Rye or Bluegrass for beautiful winter greenage.

Establishment of Emerald Zoysia Lawns By Sprigging

Establish the grade of the lawn and smooth it with a rake. Prepare the area as for a garden and apply lime if the soil test indicates its need. A soil pH of 6 to 6.5 is most desirable. Broadcast 20 to 30 pounds of a complete fertilizer such as 8-8-8 to each 1000 square feet prior to planting. Mix this into the soil by using a spade, plow, disc or Roto-Hoe. Level the area with a rake and lay off your rows six inches apart with a small plow or si-

miliar instrument. Plant the rootlets or sprigs six inches apart and approximately two inches deep, making sure that some leaves of each plant are remaining above ground level. Be sure to water each row as you plant it.

One square yard of Emerald Zoysia will plant about 1000 square feet when set on six inch centers. Newly planted lawns should be kept moist until active growth of the plants is observed. As soon as most of the plants have started spreading growth, five pounds of Nitrate of Soda per 1000 square feet should be applied to the area weekly to encourage the spread of the runners. This application should be made when the leaves are dry and it should be watered in immediately to prevent discoloration of the leaves.

Establishment of Lawns by Plugging

The use of 2-inch diameter plugs of Emerald sod instead of springs has several advantages:

- 1. There is less shock to the plants in transplanting because the roots are able to supply the leaves with more moisture immediately after planting. Thus, there is less danger of failure of the lawn to become established through insufficient moisture and less frequent watering is required during the first two weeks after planting.
- 2. Since less time is required for root establishment and more plant material is present initially, spreading will be earlier and faster.
- 3. Since plugs are better able to compete with other plants, they may be successfully used for planting directly into established bluegrass or red fescue lawns where the weeds will be controlled only by mowing.
- 4. Late summer plantings are best made with plugs because they are less subject to heaving in the North, they are less subject to loss from soil washing during the winter and because they survive better during the long winter period of no growth.

The plugs should be kept moist from in time they are cut in the nursery until they are rooted after transplanting. We sell a special hard steel plug cutter for the home owner to make accurate holes in his lawn area to fit our plugs. The plugs may be planted in rows 12 inches apart with the plugs 12 inches apart in the row. After the holes are cut they are filled with water and the Emerald plugs are placed into the holes

and stepped upon to firm the soil around the roots. The area should be watered heavily after the transplanting is completed.

The cultural practices are the same as for sprigs excepting that less frequent watering is required the first few weeks.

How To Control Weeds

It is recommended that chemical weed control should begin as soon as the grass planted and the treatment repeated every time numerous weed seedlings begin to sprout. Follow manufacturers directions as printed on each container of 2, 4-D. These treatments should not be closer than two weeks apart as injury to the grass may otherwise result. Some handweeding will be necessary if weeds are present which are not susceptible to 2, 4-D in the seedling stage or if weeds get a head start before the chemical is applied. Any hand-weeding or mechanical treatment which breaks off or uproots the runner tips of Emerald Zoysia will reduce the rapidity of coverage, therefore care in such operations is recommended.

For The Most Beautiful Lawn In The Neighborhood

Apply 15 pounds of complete fertilizer such as 8-8-8 per 1000 square feet in the spring after the last killing frost date in your locality. Topdress with 5 pounds of nitrate of soda every 6 weeks during the summer. Spread the fertilizer evenly and water it in to prevent burning the grass. Mow as needed at ½ to ¾ inch height during the summer, and use a grass catcher on the mower to remove the clippings from the lawn. During extended drough periods, best appearance will be preserved if the lawn is watered occasionally to a depth of 4 to 6 inches. Never just sprinkle lightly to cool it off.

Every three to four years the soil in the lawn area should be tested for acidity, phosphorus and potash. Recommendations obtained for correcting the deficiencies should be followed precisely.

(This information is based upon personal experience and results issued by the Georgia Agricultural Exp. Stations, U. of Ga. College of Agri., Mimeograph Series N. S. 10, dated June 1955).

NORDAN'S GRASS FARMS
"Highest Quality of Superior
Turf Grasses"
ABBEVILLE, ALABAMA

EMERALD ZOYSIA

"The Modern Miracle Lawn Grass"

Emerald Zoysia is the first released hybrid between a course textured Japanese Zoysia and a very fine textured Mascarene Zoysia. This offspring has proven superior to all other Zoysia grasses tested for turf quality. Since Emerald is a hybrid from a wide cross it is not grown from seed, but is reproduced vegetatively from sod, sprigs or plugs. The only assurance of getting genuine Emerald Zoysia is by purchasing certified stock. Most cutstanding advantages of Emerald Zoysia are:

- 1. The turf quality is unexcelled by any of our other lawn grasses. Emerald is a fine-leafed, dense growing, dark green grass that is thrillingly spongy to walk upon.
- A mature Emerald lawn is so thick and dense that weeds haven't a chance to sprout.
- 3. Emerald grows so slowly that it is no problem to keep out of flower heds, walkways and gardens.
- 4. It is possible to produce a covered lawn area in three months when planted on six inch centers.
- 5. Emerald is shade tolerant even in the Deep South where it may be found growing right up to the base of large trees.
- 6. Emerald Zoysia is more frost-tolerant than any of our other lawn grasses, therefore provides a green lawn for a much ionger sesson. It stays green the year around in some southern parts of the nation.
- 7. Emerald turf is highly resistant to wear.
- 8. The upright growth of the leaves is very slow, therefore it requires less frequent mowings.
- 9. No serious damage from diseases or insects, has been observed on Emerald Zoysla.
- 10. Emerald Zovsia produces such a dense turt that it maintains its erect body throughout the winter and lends itself to overseeding with Rye or Bluegrass for beautiful winter greenage.

Establishment of Emerald Zoysia Lawns By Sprigging

Establish the grade of the lawn and smooth it with a rake. Prepare the area as for a garden and apply time if the soil test indicates its need. A soil pH of 6 to 5.5 is most desirable. Broadcast 20 to 30 pounds of a complete fertilizer such as 8-8-8 to each 1000 square feet prior to planting. Mix this into the soil by using a spade, plow, disc or koto-Hoe. Level the area with a rake and lay off your rows six inches apart with a small plow or six inches apart with a small plow or si

miliar instrument. Plant the rootlets or sprigs six inches apart and approximately two inches deep, making sure that some leaves of each plant are remaining above ground level. Be sure to water each row as you plant it.

One square yard of Emerald Zoysia will plant about 1000 square feet when set on six inch centers Newly planted lawns should be kept moist until active growth of the plants is observed. As soon as most of the plants have started spreading growth, five pounds of Nitrate of Soda per 1000 square feet should be applied to the area weekly to encourage the spread of the runners. This application should be made when the leaves are dry and it should be watered in immediately to prevent discoloration of the leaves.

Establishment of Lawns by Flugging

The use of 2-inch diameter plugs of Emerald sod instead of springs has several advantages:

- 1. There is less shock to the plants in transplanting because the roots are able to supply the leaves with more moisture immediately after planting. Thus, there is less danger of failure of the lawn to become established through insufficient moisture and less frequent watering is required during the first two weeks after planting.
- 2. Since less time is required for root establishment and more plant material is present initially, spreading will be earlier and faster.
- 3. Since plugs are better able to compete with other plants, they may be successfully used for planting directly into established bluegrass or red fescue lawns where the weeds will be controlled only by mowing.
- 4. Late summer plantings are best made with plugs because they are less subject to heaving in the North, they are less subject to loss from soil washing during the winter and because they survive better during the long winter period of no growth.

The plugs should be kept moist from in time they are cut in the nursery until they are rooted after transplanting. We sell a special hard steel plug cutter for the home owner to make accurate holes in his lawn area to fit our plugs. The plugs may be planted in rows 12 inches apart with the plugs 12 inches apart in the row. After the holes are cut they are filled with water and the Emerald plugs are placed into the holes.

and stepped upon to firre the soil around the roots. The area should be watered heavily after the transplanting is completed.

The cultural practices are the same as for sprigs excepting that less frequent watering is required the first few weeks.

How To Control Weeds

It is recommended that chemical weed control should begin as soon as the crass planted and the treatment repeated every time numerous weed seedlings begin to sprout, Follow manufacturers directions as printed on each container of 2. 4-D. These treatments should not be closer than two weeks apart as injury to the grass may otherwise result. Some handweeding will be necessary if weeds are present which are not susceptible to 2, 4-D in the seedling stage or if weeds ger a head start before the chemical is applied. Any hand-weeding or mechanical treatment which breaks off or uproots the runner tips of Emerald Zoysia will reduce the rapidity of coverage, therefore care in such operations is recommended.

For The Most Beautiful Lawn In The Neighborhood

Apply 15 pounds of complete fertilizer such as 8-8-8 per 1000 square feet in the pring after the last killing frost date in your locality. Topdress with 5 pounds of nitrate of soda every 6 weeks during the summer. Spread the fertilizer evenly and water it in to prevent burning the grass. Mow as needed at ½ to ¾ inch height during the summer, and use a grass catcher on the mower to remove the citippings from the lawn. During extended frough periods, best appearance will be preserved if the lawn is watered occasionally to a depth of 4 to 6 inches. Never just sprinkle lightly to cool it off.

Every three to four years the soil in the lawn area should be tested for acidity, phosphorus and potash. Recommendations obtained for correcting the defleciencies should be followed precisely.

(This information is based upon personal experience and results issued by the Georgia Agricultural Exp. Stations, U. of Ga. College of Agri., Mimeograph Series N. S. 10, dated June 1955).

NORDAN'S GRASS FARMS
"Highest Quality of Superior
Turt Grosses"
ABSEVILLE, ALABAMA

Comparative Chart of Our Principle Permanent Lawn Grasses

	(1) Regions	(2) Shade	(3) Fertility	Soil	Ho₩	Mowing	(7)	(8)	(9) Wear	(10) Insect	(11) Disease	(12) Rate of	(13) Drought
GRASS	Adapted	Tolerance	Requirement	Preference	Established	Requirements	Appearance	Texture	Resistance	Damage	Trouble	Spread	Tolerance
BERMUDA				Well		Very		Extra					
South African	I, II	Poor	High	Drained	Sprigs	Frequent	Excellent	Fine	Fair	Variable	Variable	Fast	Fair
BERMUDA				Well		Very			Very	Seldom	Seldom		Very
Tiffine	1, 11, 111	Poor	High	Drained	Sprigs	Frequent	Unsurpassed	Fine	Good	Serious	Serious	Fast	Good
BERMUDA				Well		Very		Medium	Very	Seldom	Seldom		Very
Tiflawn	I, II, III	Poor	High	Drained	Sprigs	Frequent	Excellent	Fine	Good	Serious	Serious	Fast	Good
BERMUDA			And the second s	Well		Very		Medium	Very	Seldom	Seldom		Very
U-3	I, II, III	Poor	High	Drained	Sprigs	Frequent	Excellent	Fine	Good	Serious	Serious	Fast	Good
BITTER-BLUESTEM		Very		Low				Very			Seldom		Fair to
(Improved St. A.)	I	Good	Moderate	Soils	Sprigs	Frequent	Good	Coarse	Fair	Serious	Serious	Moderate	Good
CENTIPEDE					Sprigs					Seldom	Seldom		
Chinese	I, II	Medium	Low	Little	and Seed	Occasional	Good	Coarse	Fair	Serious	Serious	Moderate	Fair
ST. AUGUSTINE		Very		Low									Fair to
Common	H	Good	Moderate	Soils	Sprigs	Frequent	Good	Coarse	Fair	Serious	Serious	Moderate	Good
ZOYSIA JAPONICA					Sprigs					Seldom	Seldom		Fair to
Common	1, 11, 111	Good	Moderate	Little	and Seed	Frequent	Fair	Coarse	Good	Serious	Serious	Moderate	Good
ZOYSIA JAPONICA	201			-	2	127	THE CONTRACT	m/p/)		Seldom	Seldom		Fair to
Meyer	I, II, III	Good	Moderate	Little	Sprigg	Requent y Excellent	Excellent	Intermediate	Good	Serious	Serious	Moderate	Good
ZOYSIA MATRELLA		Very						ś	Very	Seldom	Seldom		Fair to
Common No. 13521	I, II	Good	Moderate	Little [Saries	Frequent	Excellent	Printe	Good	Serious	Serious	Slow	Good
ZOYSIA TENUFOLIA				2 × 5	111111111111111111111111111111111111111	None To	5	Extra		Seldom	Seldom		
Korean Velvet	H	Medium	Moderate	Little	Sprigs	Seldom	Excellent	·Fine	Fair	Serious	Serious	Slow	Poor
ZOYSIA				4		300	TO CO	40.	Very	None	None		,
Emerald (New Hybrid) I, II, III Good	I, II, III	Good	Moderate	Little	Sprigs	Occasional-	Unsurpassed	Fine	Good	Known	Known	Moderate	Good

We do not recommend Carpet, common Bermuda, Bahia grasses or Dichondra for lawn and turf use. Each is less desirable than the better grasses listed above, more limited in use, and more difficult to manage. Anything saved in initial establishment cost of such ground cover-will be lost in one year if adequately managed to produce presentable lawn

1. Area I is the southernmost area of the United States, including the Atlantic Coast from Charleston across the lower half of Georgia and along the Gulf of Mexico, extending inland 100 to 200 miles on through Texas. Area II extends from Noriolk southwestward through the Carolinas, lower Tennessee across into Southwestern Oklahoma on the north, and down to the Area I lines on the South. Area III covers the remainder of Oklahoma, across middle Missouri, most of Illinois, Indiana and out to the Atlantic Coast through the lower part of New York on the north, with Area II being the southern boundary. All areas have exceptions, such as mountains where the cool season grasses perform satisfactorily in the summer, and river and lake areas where winter temperatures are held up by large bodies of water.

2. Shade tolerance is extremely important in selecting a grass. Thought should be given to future shading when landscaping new lawns. All of the above grasses perform satisfactorily in full sun, but vary widely in degrees of shade that they will tolerate. Whenever possible, one grass should be used for the entire lawn.

3. Fertility requirement is that amount necessary for best performance.

4. All above grasses perform well on soils ranging from slightly acid to neutral. St. Augustine and Zoysia matrella will tolerate more salt water spray than the others.

5. Most of the recommended grasses either do not seed true to type or else seed are not commercially available. Sprigging is the surest and most economical method of establishing the better turf grasses.

6. Mowing requirement is based on that required for top performance and beauty of each particular grass. All grasses should be mowed often enough that the amount cut does not exceed the amount remaining. The finer textured grasses do best

mowed at heights of from one half to one inch while the coarser grasses should be mowed at heights of from one to two inches.

7. Appearance rating is based on a particular grass planted in an adapted area and properly maintained. If these requirements are not met, then appearance rating would be incorrect.

Texture is rated on visible leaf and stem structure.

 Wear resistance of the above grasses is satisfactory for average lawn use, but becomes extremely important in the case of institutional lawns, golf courses, athletic fields, etc.

athletic fields, etc.

10. & 11. Prospective insect and disease damage to any grass is difficult to appraise, except in the case of Common St. Augustine, which is highly susceptible to chinch bug and fungus. Most other grasses are deemed superior because they have more resistance to both insects and diseases, however all grasses are subject to damage from army worms.

12. Complete coverage from twelve inch centers on well prepared seedbeds during warm weather will vary from 8 weeks on the "fast growers" to a year or more on some of the Zoysias. Slow growth is one of the desirable characteristics of a good permanent lawn grass, and home owners will do well to reconcile themselves to spending one to two years in perfecting their lawns.

spending one to two years in perfecting their lawns.

13. Drought tolerance is an important factor that many people overlook in selecting a lawn grass. Supplemental watering is required at times for top performance on all grasses and is especially necessary during the establishing period.

NORDAN'S GRASS FARMS-"Highest Quality of Superior Turf Grasses"-Abbeville, Ala.

MORDAN'S CRASS FARMS."Highest Quality of Superior Turt Crasses"-Abbeville, Ala.

6. Mowing requirement is based on that required for top performance and beauty of each particular grass. All grasses should be mowed often enough that the amount of each particular grasses. All grasses about a powed often enough that the same particular grasses do best out does not exceed the amount remaining. The thirt textured grasses do best

Fertility requirement is that smount necessary for best performance.

All above grasses perform well on soils ranging from slightly acid to neutral.

All above grasses perform well to soils ranging thom slightly acid to neutral.

be given to future shading when landscaping new lawns. All of the above grasses beforen satisfactorily in full sun, but vary widely in degrees of shade that they will

Whenever possible, one grass should be used for the entire lawn.

are not commercially available. Sprigging is the gurest and most economical method

Most of the recommended granges either do not seed this to type or else seed

of establishing the better turt grasses.

The state of the s		CANON	THE POTTON OF	1	1	TRI MERCINO	December 10 ct. o	5111.5	0000	PUOMI	TWORK	34615DUM	Cond
NOYSIA Mendold Walter Hobertal	1 11 111		Medican	gard.		7 E 100 (6.0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	ຫ. 2	0	Very	Мопе	Моле		S
Korean Velvet)(Medium	Moderate	Little	-	Seldom	Excellent	ent'i.	Tier	200792	Serious	SJow.	P001
ALLOHUNAT AIBYON				1	5	of Sadt		EXTES	A STATE OF THE PARTY OF THE PAR	Seldom	Seldom		
Commen No. 13521	I, 11	Cooq	Moderate	Dittle 7 Spries		- Bredhant Sxurlex	Exactless !	Chimbs.	Cooq	Seldom	Seldom Serious	Wola	Fair to Good
Meyer	111.11	5000	Moderate	THIE	発展で	i whelled A the upsate i the interest in the	The House !	* Intermediate	Good	Serious	Serious.	Moderate	Good
ADINOGAL AISYOT	;	1			4	404 120	To Mark E C.		We can be a second of the seco	Seidom	Seldom		Fair to
Commen	III, III, III	Good	Moderate	Little	base bas	Етефиемі	Fair	Coarse	Good	RITOITE	Serious	Moderate	Good
ADINOGAL AISYOT					Sprigs			Abordant American Strategies in a department of the strain	And the second of the second o	Seldon	Seldom		of Tish
Common	Н	Good	Moderate	Soils	agirda	Frequent	Good	Course	Tiell	Serious.	Stroins	Moderate	Cood
SHITSURUA TE		Very		Worl		A SA CALLER OF THE SAME OF THE	A COLOR OF THE PERSON OF THE P	to region a case throughout to the second	and the second s				Fair to
Chinesa	1, 11	Medium	Low	Little	bes2 bus	Gecasional	Good	Coarse	Tish	Serious	Serious	Moderate	Tie'T
CENTIPEDE					0.07125	The state of the s	The same of the sa	Andreas and the Control of the Contr		Geldon.	Seldom		
(IA . 12 bevoram!)	~1	Cooq	Moderate	Silos	Syrigs	Frequent	್ರಂಂಥ	Coarse	7.57	Sugires	Serious	Moderate	Cooq
BITTER BLUESTEM		Very	the area of the second and the secon	Low.			The same of the sa	Very		-	Caldon	and the same of th	Nair to
D	TITI	Toc4	High	Drained	231142	Frequent	Excellent	Medium Fins	Very Good	Seldom	Seldom Serious	1285	Very Good
I Lister	3.14 418	1,001	HIST	Demist	831192	inaupar,	Excellent	Suri	Goog	Serious	Stroug	F321	Good
DATE DOK		Č		Well	1	Very			Very	Seldom			Acra
Tittine	I, II, III Poor	T00T	Migh	Drained	डिक्रां च्य	frequent	Unsurpassed	Fine	Coog	Serious	Serious	1285	Coog
ACUMERE				Well	is only only from the filling to seen such and the seen of	Very	And the second of the second o	- And the second	Very	Seldom	1	-	Very
South African	1, 11	1000	High	Drained	sgirda	Frequent	Excellent	Fire	Tight	Variable	Variable	1267	Tie?
AGUMRAG		THE RESERVE OF THE PROPERTY OF		Well		Very		Extra					
SEARIO	bestanked	Tolerance	Reduitement	Preference.	bertaildarea	Rednirements	SpharagadA	Texture	SonatalseR	Damage	Trouble	Spread	Tolerance
	840 008	ಆರಿಕ್ಕಾರಣ	F4211114W	100	Tos	MONIUM			News	pasect	Disesse	To ste A	Drought
	Sugar	(\$)	(a)	(4)	(a)	(0)	(8)	(8)	(e)	(10)	gent denn denn denn denn	(15)	(23)
ert om Lindstifferen filt i 1980 tillsam vilkelikeren men en in de skrigetillen for de		(3)	14	1 1/2 /	(27)	10.	124	Control of the Contro	n, in dans manner in administration compact distribution with a compact and co	The second secon	10/	(0)	

Comparative Chart of Our Principle Permanent Lawn Grasses

a presentable lawn. We do not recommend Carpet, common Bermuda, Babia grazses or Dichondra for la wn and turt use. Each is less desirable than the petter grazses listed above, more in land to tree and more difficult to manage. Anything saved in initial establishment cost of such ground cover-will be lost in one year it adequately managed to produce

being the southern boundary. All greas have exceptions, such as mountains where the cool season grasses perform satisfactorily in the summer, and river and isks greas the cool season grasses perform satisfactorily in the summer, and river and isks greas the cooler where winder temperatures are held up by large bodies of water.

3. Shade tolerance is expremely important in selecting a grass. Thought should extending inland 100 to 200 miles on through Texas. Area II extends from Worldik southwestern the Carolinas. Iswer Tennesses across into Southwestern Oktahoma on the north, and down to the Arca I lines on the South. Area III covers the remainder of Oklahoma, across middle Missouri, most of Illineis, Indiana and our to the adlantic Coast through the lower part of New York on the north, with Area II i. Area I is the conthermnest area of the United States, including the Atlantic Oparation Charleston across the lower half of Georgia and along the Gall of Mexico.

> 7. Appearance rating is based on a particular grazs planted in an adapted area and properly maintained. If these requirements are not met, then appearance Texture is rated on visible leaf and stem structure.

mowed at heights of from one half to one inch while the coarser grasses should

mowed at beights of from one to two inches.

but becomes extremely important in the case of incitutional lawns, golf courses, Wear resistance of the above grasses is satisfactory for average lawn use,

athletic fields, etc. 10. % It. Prospective insect and disease damage to any grass is difficult to appraise, except in the case of Common St. Augustine, which is highly suggestible to

chinch bug and finights. Most other grasses are deemed superior because they have more resistance to both inscense and diseases, however all grasses are subject to

damage from aimy worms. Complete coverage them twelve thich centers on well prepared seedbeds dur-

ang warm weather will vary from 8 weeks on the "fast growers" to a year or more on some of the Noysias. Slow growth is one of the desirable characteristics of a good spending one to two years in perfecting their lawns. 13. Diought tolerance is an important factor that many people overlook permanent lawn grass, and home owners will do well to reconcile themselves

selecting a lawn grass. Supplemental watering is required at times for top performance on all grasses and is especially necessary during the establishing period.

U-3 BERMUDAGRASS

♦ Cyondon dactylon ♦

U-3 Bermuda is a fine-bladed strain of Bermudagrass which was selected at Savannah, Georgia in 1938. It has grown in the vicinity of Washington, D. C. since then with no sign of winter injury. It has also been grown without winter-kill in the states of Washington, Nebraska and Pennsylvannia as well as in Canada.

U-3 Bermuda has its greatest use in the "crabgrass belt," which is roughly a triangular area cornered on Philadelphia, Richmond and St. Louis. It makes its best growth under the very conditions commonly associated with the worst crabgrass infestations; hot humid weather and plenty of sunlight. It is relatively indifferent to quality of soil.

U-3 provides better turf for golf course tees and fairways, putting greens, athletic fields, playgrounds, park areas and sunny lawns. Its outstanding feature is its ability to form a very wear-resistant turf that recovers rapidly. The turf is drought-resistant and can be mowed as closely as necessary to meet the demands of its use.

Insects and diseases are not yet a serious problem in growing U-3. So far, it has seldom been necessary to apply any insecticides or fungicides. An occasional attack of cutworms and sod webworms has been recorded.

U-3 Bermudagrass seed is not available, therefore it is necessary to plant it vegetatively by either of two methods: (1) plugging or (2) sprigging.

U-3 Bermuda can be planted into an existing lawn by either of these methods. It is not necessary to destroy your present lawn. A handy inexpensive

lawn plugger is now on the market to make the desired size holes to fit the purchased cut plugs. Holes are made into existing lawn area on twelve-inch centers and the cut plugs are placed into the holes and pressed firmly in place with the heel. The same process may be used in establishing U-3 onto a freshly prepared area or plugs may be hand spaced on twelve inch centers into three-inch deep rows or furrows.

The most inexpensive method of establishing U-3 is by sprigging. The U-3, solid sod less soil, is pulled apart and cut or broken into individual sprigs. The sprigs are placed into rows or furrows two-inches deep and twelve-inches apart and completely covered with soil every twelve-inches in the row.

When planting into an existing lawn, the sod is opened with an edging tool, or by similiar means and the sprigs are placed into the trenches in the same manner as above. Soil should be firmed in place by tampering, rolling, watering or with foot pressure.

Rate of coverage depends primarily on the distance apart the plugs or sprigs are placed. On newly prepared seed beds, plugs placed at one-foot intervals will require from 6 to 8 weeks to form solid turf; plugs placed at two-foot intervals may form solid sod within one growing season.

Plugs placed at one-foot intervals into existing lawns will require one growing season to form a turf; two-foot centers will require a second growing season.

U-3 sprigged on eight-inch centers into a well prepared new lawn area and

kept free of competitive weeds will produce solid sod in six weeks; on 8 to 12 inch centers from 8 to 12 weeks and on 18 to 24 - inch centers one complete growing season.

U-3 sprigged into existing lawns on 8 to 12-inch centers will produce turf in one growing season; 18 to 24 - inch centers will require a second growing season.

The newly-planted area should be kept moist until the grass is well established. This generally requires watering once a day for the first week, then several times per week for the next few weeks.

U-3 Bermudagrass should be fertilized at the rate of 40 to 50 pounds of a 10-6-4 fertilizer per 1,000 square feet the first year. One-third should be applied at time of planting, one-third three weeks after planting and one-third six weeks after planting.

Overseeding for winter color may be done in the fall with a permanent coolseason grass mixture or with a temporary grass such as redtop or domestic ryegrass.

We wish you success in establishing a U-3 lawn. We stand ready to supply more plugs or sprigs as you want to expand your plantings. U-3 may be planted successfully all summer long, even during the hottest weather.

NORDAN'S GRASS FARMS

"Growers of Superior Turf Grasses"

Abbeville, Alabama

U-3 BERMUDAGRASS

🛊 Cyondon daetylon 🎕

U-3 Bermuda is a fine-bladed strain of Bermudagrass which was selected at Savannah, Georgia in 1938. It has grown in the vicinity of Washington, D. C. since then with no sign of winter injury. It has also been grown without winter-kill in the states of Washington, Nebraska and Pennsylvannia as well as in Canada.

U-3 Bermuda has its greatest use in the "crabgrass belt," which is roughly a triangular area cornered on Philadelphia, Richmond and St. Louis. It makes its best growth under the very conditions commonly associated with the worst crabgrass infestations: hot humid weather and plenty of sunlight. It is relatively indifferent to quality of soil.

U-3 provides better turf for golf course tees and fairways, putting greens, athletic fields, playgrounds, park areas and sunny lawns. Its outstanding feature is its ability to form a very wear-resistant turf that recovers rapidly. The turf is drought-resistant and can be moved as closely as necessary to meet the demands of its use.

Insects and diseases are not yet a serious problem in growing U-3. So far, it has seldom been necessary to apply any insecticides or fungicides. An occasional attack of cutworms and sod webworms has been recorded.

U-3 Bermudagrass seed is not available, therefore it is necessary to plant it vegetatively by either of two methods: (1) plugging or (2) sprigging.

U-3 Bermuda can be planted into an existing lawn by either of these methods. It is not necessary to destroy your present lawn. A handy inexpensive

lawn plugger is now on the marker to make the desired size holes to fit the purchased cut plugs. Holes are made into existing lawn area on twelve-inch centers and the cut plugs are placed into the holes and pressed firmly in place with the heel. The same process may be used in establishing U-3 onto a freshly prepared area or plugs may be hand spaced on twelve inch centers into three-inch deep rows or furrows.

The most inexpensive method of establishing U-3 is by sprigging. The U-3, solid sod less soil, is pulled apart and cut or broken into individual sprigs. The sprigs are placed into rows or furrows two-inches deep and twelve-inches apart and completely covered with soil every twelve-inches in the row.

When planting into an existing lawn, the sod is opened with an edging tool, or by similiar means and the sprigs are placed into the trenches in the same manner as above. Soil should be firmed in place by tampering, rolling, watering or with foot pressure.

Rate of coverage depends primarily on the distance apart the plugs or sprigs are placed. On newly prepared seed beds, plugs placed at one-foot intervals will require from 6 to 8 weeks to form solid turf; plugs placed at two-foot intervals may form solid sod within one growing season.

Plugs placed at one-foot intervals into existing lawns will require one growing season to form a turf; two-foot centers will require a second growing season.

U-3 sprigged on eight-inch centers into a well prepared new lawn area and

kept free of competitive weeds will produce solid sod in six weeks; on 8 to 12 meh centers from 8 to 12 weeks and on 18 to 24 - inch centers one complete growing season.

U-3 sprigged into existing lawns on 8 to 12-inch centers will produce turf in one growing season; 18 to 24 - inch centers will require a second growing season

The newly-planted area should be kept moist until the grass is well established. This generally requires watering once a day for the first week, then several times per week for the next few weeks.

U-3 Bermudagrass should be fertilized at the rate of 40 to 50 pounds of a 10-6-4 fertilizer per 1,000 square feet the first year. One-third should be applied at time of planting, one-third three weeks after planting and onethird six weeks after planting.

Overseeding for winter color may be done in the fall with the permanent colsection grass mixture or with a temporary grass such as redtop or domestic ryegrass.

We wish you success in establishing a U-3 lawn. We stand ready to supply more plugs or sprigs as you want to expand your plantings. U-3 may be planted successfully all summer long, even during the hottest weather.

NORDAN'S GRASS FARMS

"Growers of Superior Turf Grasses"

Abbeville, Alabama

Meyer *₹-52* Zoysia Japonica

H It's Permanent

* No More Replanting

* Dark Green All Summer Long

Meyer (Z-52) Zoysia is one of the new improved turf grasses which gives hope to the homecomer who desperately wants a good lawn free from Crabgrass. This was developed, tested and released as a superior turf grass for lawns and specialized turf areas by the United States Department of Agriculture and U. S. Golf Association, Green Section. Meyer (Z-52) Zoysia is a revelation to people who walk upon it for the first time. Once established, it produces a dense resiliant and weed free turf. It is noted for its "dream come true" dark green which is so essential for maximum lawn beauty.

Yes, a mature Meyer Zoysia lawn:

- 1-is permanent.
- 2-is constantly improving with age.
- 3—is free from Crabgrass and other summer weeds.
- 4-is free from diseases and insects.
- 5—is extremely resistant to heat and drough.
- 6-thrives in sun or shade.
- 7—is highly wear resistant tests have proven that it will withstand more wear than other grasses; excessive play and recreation will not wear paths through it.
- 8—is a beautiful dark green turf that remains dense and resiliant throughout the year.
- 9—is readily adaptive to overseeding with cool-season grasses for midwinter greenage.
- 10—is excellent for banks, terraces and erosion control.
- 11—requires less frequent mowings—thrives under close mowing.
- 12—stands neglect without tolerating invading weeds.
- 13—is an evident luxury, setting off the house and garden more effectively than a Cadillac or Lincoln parked in the driveway.

- 14—saves time by forcing the use of the power mower, an effeciency step never again to be surrendered.
- 15—is a beautiful green color when most cool-weather grasses have turned brown.

All of this and more too—Meyer is completely indifferent to soil composition—will grow in poor soils — sandy soils— clay soils— or "so called" rock beds.

Exclusion of noxious weeds, especially Crabgrass, is now possible with this new superior turf grass. Meyer (Z-52) Zoysia is most aggressive and dominant during the crabgrass season. Once established, this grass is highly resistant to weed invasion due to the tight interwoven mass of turf formed.

Meyer (Z-52) Zoysia forms a beautiful fine textured lawn. It may be cut at a height of from three-eights of an inch to as high as four inches. The backbone of Meyer (Z-52) Zoysia is made up of short stems and tough interwoven stolons. These and many underground rhizomes with their dense mat of roots greatly enhance the wearing and load supporting quality of the sod. Combine this with the upright growth of the blades and one can well imagine the mat of cushiony turf unequalled for wear resistance and resilence.

Meyer (Z-52) Zoysia has withstood severe winter conditions as far north as Canada and thrives in the extreme heat of Southern Florida. It becomes dormant and loses its green color when freezing weather begins in the fall, however it never loses its dense turf quality. It regains its dark green color in the spring when the weather becomes warm. This presumed disadvantage of color loss is easily overcome by the seeding of cool season grasses, such as Merion Bluegrass into the Zoysia turf in the fall of the year.

There is evidence that cool - season grasses thrive better in a Zoysia base than they do alone. The Zoysia base

provides a cooler soil during the summer months and carries the cool-season grass through its most dangerous period in good shape.

Meyer (Z-52) Zoysia is eagerly awaited by home owners and turf workers throughout the country who have regularly experienced "loves labor lost."

Seed of this superior strain of Zoysia is not available. Planting must be done vegetatively. The simplest and easiest method of establishing a Meyer Zoysia lawn is by planting plugs. It is not necessary to work up your present lawn and start with a newly prepared seed bed. Meyer (Z-52) Zoysia can easily be planted into an existing lawn with the aid of a special inexpensive plugger-transplanters now on the market.

Zoysia plugs, set on one foot centers into your existing lawn will afford coverage within a period of two growing seasons. In a newly prepared seed bed the time is cut into half, one year from one foot centers. If anything, these figures are conservative.

Sprigging is the more inexpensive method used for establishing Meyer onto a new lawn area. Solid Meyer sod minus soil is shipped for this purpose. The sod is divided into individual plants of one root system and one leaf cluster and planted on twelve inch centers. Meyer may be forced to complete coverage in less than four months by using this method and by giving proper care.

We strongly recommend that part of your trial planting be made in your most troublesome lawn area to prove that this is truly a superior turf grass. The relatively high cost of the initial planting material is in reality an economy because the Meyer not only improves the planted area but will also serve as a nursery for plugging to set additional turf areas.

NORDAN'S GRASS FARMS

"Highest Quality of Superior Turf Grasses"

ABBEVILLE, ALABAMA

ZOYSIA MATRELLA

"King of the Lawn Grasses"

The New Miracle Grass

Thickest Carpet of Any Grass

* The Toughest Lawn Grass

Yes, a mature Zoysia Matrella lawn:

1—is the most beautiful of all lawns.

2-is permanent.

3-is constantly improving.

4-is free from weeds.

5—is free from disease and insect enemies (no more chinch bug trouble).

6-is drough resistant.

7-thrives in shade and sun.

8—stands hard use better than any other grass.

9—stands neglect without tolerating invading weeds.

10—is an evident luxury often referred to as the "rich man's grass".

11—saves time by forcing the use of a power mower — an efficiency step never again to be surrendered.

Zoysia matrellas are native to the Orient where they are used for turf forage and fuel. They were introduced into this country in 1911 for observation and experimentation. Recently, they have been acclaimed as the very best all purpose lawn grass for the South. They grow well in all sections of the Southern part of the United States but are not recommended in areas north of Washington, D. C.

Zoysia matrellas have fine leaves, dark green in color and produce a dense sod which gives one the sensation of walking on a deep piled Persian carpet. One of their most valued characteristics is, after reaching maturity, their ability to eradicate all weeds. They are one of the earliest grasses to begin growth in the spring and one of the last to go dormant in the fall—remaining green much longer than Bermuda.

Matrellas grow to a height of only three or four inches and require few and infrequent mowings. Because of their slow growth, they need trimming from around flower beds only once yearly. Matrellas flourish under the hot sun and thrive under more shade

than any grasses tested at the Alabama Experiment Station. They do well under oaks, elms, pines, walnuts, cedars and most other trees.

Zoysia matrellas are noted for their hardy qualities being tough enough to withstand the continued play of children. They flourish and make a beautiful carpet in poor soil such as sand and clay as well as in areas favored with selected top soil. Zoysias should never be top-dressed with soil after planting.

Since seeds are not available, matrellas must be established vegetatively from sprigs or plugs set on one-foot centers. Complete coverage may be had in less than one growing season if irrigation, fertilization and weed control are practiced.

From our experience, early summer plantings produce the best results; although, many of our customers have reported sprigging and plugging Zoysias every month in the year with excellent results.

WHAT THE EXPERTS SAY ABOUT ZOYSIA MATRELLA LAWN GRASSES

AUBURN—"A dark green fine leaved lawn grass that will grow both in the sun and shade has been the goal of lawn research workers in the South for years. Zoysia matrella, a grass with these characteristics has been introduced from the Orient by the Department of Agriculture. Zoysia matrella produces a dense sod that feels like a rug. Unlike bermuda, it does not get weedy." (Dr. D. G. Sturkie, Agronomist, A.P.I.)

"ZOYSIA MATRELLA grows in pure red clay beneath pine trees where plantings of other grasses had completely failed; it grows in sand on the seashore where other grasses haven't a chance; it thrives in poor soil and flourishes in good soil. It is wellnamed 'Miracle Grass'". (W. Elbridge Freeborn, Garden Editor, The Birmingham News-Age Herald)

MONTGOMERY — "Walking on the capital lawn nowadays is like walking on a persian rug. That's because of 'Zoysia matrella . . . not only does matrella (the toughest grass ever grown) stand up under shade, wetness or long dry spells, but it kills other grasses.

"During the war, according to Joe H. Abercrombie, capital grounds land-scaper, air pilots found that they could land their heavy planes on turfs planted with Zoysia matrella without breaking through the ground no matter if it were wet or dry.

"In parts of the South it is just as green in the Winter as in Summer—and also holds its rich green hue during Summer droughs." (International News Service)

BELTSVILLE, MD.—"In the United States, no disease has as yet been observed and identified on Zoysia matrella. It is remarkably free from insect parasites." (Dr. Ian Forbes, Jr.—Dr. Marvin Ferguson, Research Agronomists, USGA & USDA)

"ZOYSIA MATRELLA is by far the best known and best dressed of the Zoysias . . . The spongy thick turf of matrella is one of its most desirable and exclusive features . . . No other lawn grass can equal its adaptability to various types of soil and climate." (W. W. Nordan, "Green Thumb")

"ZOYSIA MATRELLA may be set with good results at any time of the year. If plenty of water is available during the dry Summer months, and a light covering of pine needles or leaves is used for the December-January settings, there is little choice between one month and another." (M. L. Beck of Auburn)

"ZOYSIA MATRELLA is free from all attacks by 'chinch bugs' and Japanese bettle grubs have never been known to invade it, no doubt because the roots are too tough. Once well established, Zoysia needs very little food and much less water than ordinary grasses." (C. W. Baker "Home Garden")

ALLES IN THE FIRST OF

As ROYAL BUILDING BUILDING